

Establishment of gene transfer and gene silencing methods in a desiccation-tolerant cell line, Pv11

Sogame Y., Okada J., Kikuta S., Miyata Y., Cornette R., Gusev O., Kikawada T.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2016, Springer Japan. Larvae of the African midge *Polypedilum vanderplanki* show extreme desiccation tolerance, known as anhydrobiosis. Recently, the cultured cell line Pv11 was derived from this species; Pv11 cells can be preserved in the dry state for over 6 months and retain their proliferation potential. Here, we attempted to expand the use of Pv11 cells as a model to investigate the mechanisms underlying anhydrobiosis in *P. vanderplanki*. A newly developed vector comprising a constitutive promoter for the *PvGapdh* gene allowed the expression of exogenous proteins in Pv11 cells. Using this vector, a stable Pv11 cell line expressing green fluorescence protein (GFP) was established and retained desiccation tolerance. Gene silencing with GFP-specific siRNAs significantly suppressed GFP expression to approximately 7.5–34.6% of that in the non-siRNA-transfected GFP stable line. Establishment of these functional assays will enable Pv11 cells to be utilized as an effective tool to investigate the molecular mechanisms underlying anhydrobiosis.

<http://dx.doi.org/10.1007/s00792-016-0880-4>

Keywords

Anhydrobiosis, Desiccation tolerant cells, Gene expression system, RNAi

References

- [1] Cornette R, Kanamori Y, Watanabe M et al (2010) Identification of anhydrobiosis-related genes from an expressed sequence tag database in the cryptobiotic midge *Polypedilum vanderplanki* (Diptera; Chironomidae). *J Biol Chem* 285:35889–35899. doi:10.1074/jbc.M110.150623
- [2] Ducrest AL, Amacker M, Lingner J, Nabholz M (2002) Detection of promoter activity by flow cytometric analysis of GFP reporter expression. *Nucleic Acids Res* 30:e65. doi:10.1093/nar/gnf064
- [3] Erkut C, Penkov S, Khesbak H et al (2011) Trehalose renders the dauer larva of *Caenorhabditis elegans* resistant to extreme desiccation. *Curr Biol* 21:1331–1336. doi:10.1016/j.cub.2011.06.064
- [4] Gusev O, Suetsugu Y, Cornette R et al (2014) Comparative genome sequencing reveals genomic signature of extreme desiccation tolerance in the anhydrobiotic midge. *Nat Commun* 5:4784. doi:10.1038/ncomms5784
- [5] Hatanaka R, Hagiwara-Komoda Y, Furuki T et al (2013) An abundant LEA protein in the anhydrobiotic midge, PvLEA4, acts as a molecular shield by limiting growth of aggregating protein particles. *Insect Biochem Mol Biol* 43:1055–1067. doi:10.1016/j.ibmb.2013.08.004
- [6] Kikawada T, Nakanahara Y, Kanamori Y et al (2006) Dehydration-induced expression of LEA proteins in an anhydrobiotic chironomid. *Biochem Biophys Res Commun* 348:56–61. doi:10.1016/j.bbrc.2006.07.003
- [7] Li S, Chakraborty N, Borcar A, Menze MA, Toner M, Hand SC (2012) Late embryogenesis abundant proteins protect human hepatoma cells during acute desiccation. *Proc Natl Acad Sci USA* 109:20859–20864. doi:10.1073/pnas.1214893109

- [8] Naito Y, Yoshimura J, Morishita S, Ui-Tei K (2009) siDirect 2.0: updated software for designing functional siRNA with reduced seed-dependent off-target effect. *BMC Bioinform* 10:392. doi:10.1186/1471-2105-10-392
- [9] Nakahara Y, Imanishi S, Mitsumasu K et al (2010) Cells from an anhydrobiotic chironomid survive almost complete desiccation. *Cryobiology* 60:138–146. doi:10.1016/j.cryobiol.2009.10.004
- [10] Okada J, Kikuta S, Gusev O et al (2015) Construction of optimized CRISPR/Cas system to reveal the mechanisms of anhydrobiosis in the Sleeping Chironomid. *Cryobio Cryotech* 61:69–73
- [11] Sakurai M, Furuki T, Akao K et al (2008) Vitrification is essential for anhydrobiosis in an African chironomid, *Polypedilum vanderplanki*. *PNAS* 105:5093–5098. doi:10.1073/pnas.0706197105
- [12] Shimizu T, Kanamori Y, Furuki T et al (2010) Dessication-induced structuralization and glass formation of group 3 Late Embryogenesis Abundant protein model peptides. *Biochemistry* 49:1093–1104. doi:10.1021/bi901745f
- [13] Ui-Tei K, Naito Y, Takahashi F et al (2004) Guidelines for the selection of highly effective siRNA sequences for mammalian and chick RNA interference. *Nucleic Acids Res* 32:936–948. doi:10.1093/nar/gkh247
- [14] Watanabe M (2006) Anhydrobiosis in invertebrates. *Appl Entomol Zool* 41:15–31
- [15] Watanabe M, Kikawada T, Okuda T (2003) Increase of internal ion concentration triggers trehalose synthesis associated with cryptobiosis in larvae of *Polypedilum vanderplanki*. *J Exp Biol* 206:2281–2286
- [16] Watanabe K, Imanishi S, Akiduki G, Cornette R, Okuda T (2016) Air-dried cells from the anhydrobiotic insect, *Polypedilum vanderplanki*, can survive long term preservation at room temperature and retain proliferation potential after rehydration. *Cryobiology* 73:93–98. doi:10.1016/j.cryobiol.2016.05.006